


Calidris pusilla Semipalmated Sandpiper

Subspecies/Population	Breeding range	Wintering, or core non-breeding range	Afr	Eu	Asia	Oc	Neo	NA	Estimate	Source	Trend	Source	1% level	Notes
Alaska (br)	W & N Alaska	<i>Pacific Coast Mexico to S Peru</i>					•	•	2,000,000–2,500,000	M42	DEC	M42		M42 presents an estimate for the 3 populations combined of 2,000,000–2,500,000 and recommends use of a midpoint of 2,250,000.
E North Canada (br)	E North Canada	<i>Atlantic Coast Yucatan & W Indies to C Argentina</i>					•	•						
W North Canada (br)	W North Canada	<i>Atlantic Coast Yucatan & W Indies to C Argentina</i>					•	•						

Calidris mauri Western Sandpiper

Subspecies/Population	Breeding range	Wintering, or core non-breeding range	Afr	Eu	Asia	Oc	Neo	NA	Estimate	Source	Trend	Source	1% level	Notes
Alaska, Chukotskiy (br)	W & N Alaska, E Chukotskiy Peninsula	<i>Coasts N Washington–S Peru, E USA, Caribbean to Suriname</i>					•	•	3,500,000	M41	DEC?	M42	20,000	For populations over 2 million birds, Ramsar Convention criterion 5 (20,000 or more waterbirds) applies.

Calidris ruficollis Red-necked Stint

Subspecies/Population	Breeding range	Wintering, or core non-breeding range	Afr	Eu	Asia	Oc	Neo	NA	Estimate	Source	Trend	Source	1% level	Notes
NE Siberia (br)	N Siberia E of C & E Taymyr S to N Kamchatka; sporadic W & N Alaska	<i>E India, Sri Lanka, through SE & E Asia to Australasia</i>					•	•	315,000	B9			3,200	Increasing in Australia (S42) but decreasing in Japan and Korea (M32), suggesting possibility of a shift in wintering range.

Calidris minuta Little Stint

Subspecies/Population	Breeding range	Wintering, or core non-breeding range	Afr	Eu	Asia	Oc	Neo	NA	Estimate	Source	Trend	Source	1% level	Notes
Europe & West Africa (non-br)	N Scandinavia, N European Russia	<i>W Europe, Black Sea, Mediterranean, N & W Africa</i>					•	•	200,000	D18 S43	DEC?	S43	2,000	
SW Asia, E & S Africa (non-br)	N West Siberia	<i>Caspian region, SW Asia, East & Southern Africa</i>					•	•	1,000,000	P10			10,000	D18 considers this estimate to be too low.
S Asia (non-br)	NW & NC Siberia to New Siberian Is & R Yana	<i>Indian Subcontinent, Sri Lanka, Myanmar</i>					•		200,000–300,000	B6			2,500	B6: Indian population estimated at 200,000.

Calidris temminckii Temminck's Stint

Subspecies/Population	Breeding range	Wintering, or core non-breeding range	Afr	Eu	Asia	Oc	Neo	NA	Estimate	Source	Trend	Source	1% level	Notes
Europe, W Africa (non-br)	Scandinavia, NW Russia	<i>Sub-Saharan W Africa via Black Sea, Mediterranean</i>					•	•	39,000–80,000	S43			600	H25: No evidence of a population trend among migrants in Sweden, but possibility of decline in Finland.
SW Asia, E Africa (non-br)	NE Europe, W Siberia	<i>NE & E Africa S to Kenya, Burundi, via SW Asia</i>					•	•	D/E	S18				
S Asia (non-br)	N Siberia	<i>S Asia</i>					•		C	B6			1,000	B6: Indian population estimated at <50,000.
E & SE Asia (non-br)	N Siberia	<i>Indochina, S China, Taiwan, Malay Peninsula, Borneo</i>					•		B/C	B9			1,000	